



**ENVIRONMENTAL
HEALTH
PERSPECTIVES**

ehponline.org

**Urinary Biomarkers of Prenatal
Atrazine Exposure and Adverse Birth
Outcomes in the PELAGIE Birth Cohort**

**Cécile Chevrier, Gwendolina Limon, Christine Monfort,
Florence Rouget, Ronan Garlantézec, Claire Petit,
Gaël Durand, and Sylvaine Cordier**

**doi: 10.1289/ehp.1002775 (available at <http://dx.doi.org/>)
Online 2 March 2011**



NIEHS
National Institute of
Environmental Health Sciences

National Institutes of Health
U.S. Department of Health and Human Services

Urinary Biomarkers of Prenatal Atrazine Exposure and Adverse Birth Outcomes in the PELAGIE Birth Cohort

Cécile Chevrier (1), Gwendolina Limon (2), Christine Monfort (1), Florence Rouget (1,3),
Ronan Garlantézec (1,4), Claire Petit (1), Gaël Durand (2), Sylvaine Cordier (1)

(1) Inserm, U625, F-35000 Rennes, France; University of Rennes I, GERHM, IFR140, F-35000 Rennes, France

(2) IDHESA, F-29000 Plouzané, France

(3) " Bien naître en Ille-et-Vilaine " Perinatal network, Rennes, France

(4) Public Health department, CHU Brest, Brest, France

Corresponding author

Cécile Chevrier

Inserm U625, University of Rennes 1, F-35000, Rennes, France

Tel: +33(0)223236126

Fax: +33(0)223235055

cecile.chevrier@rennes.inserm.fr

Acknowledgments and grant supports

We are grateful to the participants and the staff of maternity hospitals and clinics. Thanks to N.Costet, D.Marchand, L.Multigner, N.Bonvallot, V.Villalon and J.A.Cahn. This study is funded by the National Research Agency and the Regional Council of Brittany.

Competing financial interest declaration

The authors declare they have no competing financial interests.

Short running head

Atrazine Exposure and Birth Outcomes

Key words

Atrazine, environmental exposure, fetal growth, herbicides

Abbreviations

BMI: Body mass index

FGR: Fetal growth restriction

SGA: Small for gestational age

SHC: Small head circumference for sex and gestational age

LOD: Limit of detection

LOQ: Limit of quantification

Abstract

Background: Despite evidence of atrazine toxicity on developing organisms from experimental studies, only few studies – and fewer epidemiological investigations – have examined the potential effects of prenatal exposure.

Objectives: We sought to assess the association between adverse birth outcomes and urinary biomarkers of prenatal atrazine exposure, while taking into account exposures to other herbicides used on corn crops (simazine, alachlor, metolachlor, and acetochlor).

Methods: This study used a case-cohort design nested in a prospective birth cohort conducted in the Brittany region from 2002 through 2006. It collected maternal urine samples to examine pesticide exposure biomarkers before the 19th week of gestation.

Results: Quantifiable levels of atrazine or atrazine mercapturate were found in urine samples from 5.5% of 579 pregnant women, and dealkylated and hydroxylated triazine metabolites were identified in 20% and 40% of samples, respectively. The presence versus absence of quantifiable levels of atrazine or a specific atrazine metabolite was associated with fetal growth restriction (OR=1.5; 95% CI: 1.0-2.2) and small head circumference for sex and gestational age (OR=1.7; 95% CI: 1.0-2.7). Associations with major congenital anomalies were not evident with atrazine or its specific metabolites. Head circumference was inversely associated with the presence of quantifiable urinary metolachlor.

Conclusions: This study is the first to assess associations of birth outcomes with multiple urinary biomarkers of exposure to triazine and chloroacetanilide herbicides. Evidence of associations with adverse birth outcomes raises particular concerns for countries where atrazine is still in use.

Introduction

Atrazine is a triazine herbicide used to control broadleaf and grassy weeds in crops, mainly corn and sorghum. Although the European Union banned atrazine in 2001 because it is a ubiquitous water contaminant, it is still used in more than 70 countries, including the United States, Brazil, Argentina, Mexico, and China (LeBaron et al. 2008). In 2001, agricultural pesticide use in France was among the highest in Europe, averaging 3.4 kilograms per hectare of agricultural area (Benoît et al. 2005). The intensive regional agricultural activities due to pig, poultry, and dairy farming in the Brittany region (northwestern France) make pesticide exposure a critical issue. Corn crops, especially animal feed corn, are common there. In 2001, experts estimated that 200 tons of atrazine was applied to 70% of the corn grown in Brittany, for an average of 0.620 kilograms of atrazine per hectare (AGRESTE Bretagne 2001). In 1997, the Ministry of Agriculture limited atrazine to 1 kg/ha/year; in 2002, it banned the sale of numerous products containing atrazine, and in October 2003, finally banned atrazine use. The primary replacements have been alachlor and acetochlor, both chloroacetanilide herbicides.

Once released into the environment, atrazine is transformed over time by various chemical, photochemical, and biologically mediated reactions into other compounds, called degradates or metabolites. Water contamination by atrazine and its degradates, including hydroxyatrazine and desethylatrazine, has decreased in Brittany since 2003, but many surveys still show environmental contamination in the region. Testing indicates that 27% of water samples from Breton rivers had atrazine levels $> 0.1 \mu\text{g/L}$ in 2003 compared with 10% of samples in 2004, and only 1% in 2007, while atrazine degradates were detected in 76% of water samples in 2003 and 20% in 2007 (Réseau CORPEP 2008). Atrazine residues were also detected in surface and underground sources of human drinking water (Marchand 2005). In 2006 the

maximum measured surface water concentration was 0.38 µg/L for atrazine and 0.37 µg/L for hydroxyatrazine, and underground levels 0.14 µg/L for atrazine, 0.38 µg/L for hydroxyatrazine, and 0.32 µg/L for desethylatrazine. Finally, in drinking water of Brittany in 2005, when the regulatory limits of pesticides (0.1 µg/L) were exceeded, it was mainly due to desethylatrazine (67%) and atrazine (17%) residues (Marchand 2005).

Some epidemiological studies have looked at the potential impact of prenatal exposure to atrazine or its environmental degradates on pregnancy outcomes in the general population. Munger et al. (1997) collected data from 1984 through 1990 and observed higher rates of babies born small-for-gestational-age (SGA) in Iowa communities with higher atrazine levels in their drinking water. In Brittany, Villanueva et al. (2005) found that the risk of SGA increased with the number of months of the third trimester of pregnancy occurring from May to September, months with the highest atrazine levels in drinking water there from 1990 through 1998. Ochoa-Acuña et al. (2009) found a statistically significant increase in SGA prevalence associated with atrazine in drinking water in Indiana from 1993 through 2007. In Washington State, Waller et al. (2010) observed a higher risk of gastroschisis from 2001 through 2006 among babies of mothers who lived closer to sites with high atrazine concentrations. Three other studies related high incidences of birth defects to counties with high triazine usage in New York from 1995 through 1999 (Davis 2005), to the highest mean atrazine levels in surface water in Indiana from 1990 through 2002 (Mattix et al. 2007), and to the annual April-July peak of atrazine contamination of surface water (based on 1996-2002 US data, Winchester et al. 2009). Most of these studies were retrospective and relied on ecological exposure assessment. Co-occurring chemicals make it impossible to determine whether associations are specific to atrazine exposure in the general population.

Our aim is to assess the association between adverse birth outcomes and urinary biomarkers of prenatal atrazine exposure, taking into account other herbicides potentially used on corn,

including simazine, alachlor, metolachlor, and acetochlor. This study is based on a prospective population-based cohort in the Brittany region —the PELAGIE cohort — that collected urine samples from pregnant women between 2002 and 2006 to assess pesticide exposure biomarkers before the 19th week of gestation.

Materials and Methods

The PELAGIE cohort

The PELAGIE cohort included 3421 pregnant women from Brittany from 2002 through February 2006. Enrollment started in the district of Ille-et-Vilaine, continued in Finistère, and ended in Côtes d'Armor. Gynecologists, obstetricians, and ultrasonographers recruited women during consultations in early pregnancy, informing them about the nature of the study and asking them to participate (after providing written consent). Women were enrolled before the 19th week of gestation after providing written informed consent and completing a questionnaire at home concerning family social and demographic characteristics, diet and lifestyle. Women were asked to return the questionnaire by mail, along with a first-morning-void urine sample that they collected and transferred into two vials containing nitric acid (to inhibit bacterial proliferation). Samples were mailed to the study laboratory (Inserm U625) in a pre-stamped package at ambient temperature, with routine delivery taking from 1 to 3 days. Upon receipt, samples were frozen and stored at -20°C. Midwives and pediatricians at the maternity units provided study staff with medical information about the pregnancy, delivery, and neonatal health for 3399 women (99.4% of the cohort; the other 22 women were lost to follow-up). Birth weight, length, and head circumference were obtained from hospital records by pediatricians. Gestational age was based on both date of the last menstrual period and ultrasound examinations. The appropriate national ethics committees approved the study.

Outcome definition

Liveborn singleton children of the cohort without major congenital anomalies were classified as fetal growth restriction (FGR) if their birth weight was below the 5th percentile of the distribution of expected birth weight modeled within the cohort according to gestational age (third-degree polynomial), sex, parity (third-degree polynomial), maternal weight (third-degree polynomial), height, and age (second-degree polynomial) (Mamelle et al. 2001). Small head circumference (SHC) was defined by a head circumference below the 5th percentile of the birth head circumference distribution for a given gestational age and sex, according to French reference curves (Mamelle et al. 1996). Major congenital malformations in liveborn infants were diagnosed by hospital staff pediatricians at the maternity unit, based on findings at the clinical examination at birth. In addition, pediatricians were asked specifically about male genital anomalies (hypospadias, undescended testis, and micropenis). Until 2 years after birth, diagnosis validation by pediatric surgeons was obtained for half of these male genital anomalies (n=12/26). For detailed information on the classification of congenital malformations in this cohort, see Garlantézec et al. (2009).

Data on corn agricultural activities and atrazine water contamination

Data on agricultural activities were collected for each mother's municipality of residence at enrollment from the national agricultural census conducted in 2000; specifically, we recorded the proportion of the municipality's area used for corn crops. Drinking water contamination by atrazine was routinely assessed until its year of ban by the office of Social and Sanitary Affairs of Brittany. We collected these data from 2000 to 2002 for municipalities in Ille-et-Vilaine, and used them to estimate the average atrazine level in water during the first trimester of pregnancy for participants residing in these locations. Exposure to atrazine via tap water

was then estimated by multiplying the average concentrations by the tap water consumption reported in the questionnaire.

Case-cohort design

The present study was based on a case-cohort design (Ahrens and Pigeot 2004) that included 601 children randomly selected from the entire liveborn singleton cohort (18%), referred to as the subcohort, and groups with adverse birth outcomes, including congenital anomalies (88 major anomalies, 26 male genital anomalies), FGR (n=180), and SHC (n=105), representing all of the cohort members with these outcomes.

Pesticide determination in urine samples

Pesticides were measured in urine samples from 579 women in the subcohort (96%), all mothers of babies with major congenital anomalies or male genital anomalies, 178 mothers of children with FGR, and 103 mothers with children with SHC. First-morning-void urine samples, which are very likely to contain recent traces of exposure to the nonpersistent pesticides we targeted (Barr et al. 2005), were used to determine the presence of 12 triazine compounds: atrazine and its glutathione-derived metabolite atrazine mercapturate, simazine and its metabolite simazine mercapturate, three dealkylated triazine metabolites that are likely to be formed in water systems (desethylatrazine, desisopropyl atrazine, and 2-chlorodiaminoatrazine), and five hydroxylated triazine metabolites, mostly formed in plants (hydroxyatrazine, hydroxysimazine, hydroxydesethylatrazine, hydroxy-desisopropyl atrazine, and hydroxy-2-chlorodiaminoatrazine (ammeline)). The chloroacetanilide herbicides alachlor, metolachlor, and acetolachlor were also measured, together with 2,6-diethylaniline, an alachlor metabolite.

Chemical analyses were performed by the IDHESA Institute (Plouzané, Finistère) on urine samples (maximum: 10 mL) with liquid chromatography and triple quadrupole mass spectrometry (LC/MSMS) after solid phase extraction (SPE). Triazine and chloroacetanilide herbicides were analyzed simultaneously. Reference standards for pesticides and their metabolites were purchased from Dr. Ehrenstorfer (Augsburg, Germany) and from Promochem (Teddington, UK). Other purchased chemicals included: LC-MS grade acetonitrile and methanol from Fisher Scientific (Loughborough, Leicestershire, UK) (>99%), analytical grade formic acid from Baker (Deventer, Netherlands) (98%), and gas from Air Liquide (Paris, France) (>99%). Standard solutions were prepared in methanol or acetonitrile and stored at -20 °C. Simazine-d10 was used as internal standard for extraction and detection controls. Calibration standards were prepared by adding appropriate working standard solutions to 10-mL fresh samples of pesticide-free human urine before extraction to obtain concentrations in the calibration range. Isotope-labeled standards and surrogates were added to calibration standards to obtain a final concentration of 1 µg/L. The urine samples were thawed and shaken. Preconcentration and extraction (5 ml) was performed with an on-line SPE high volume symbiosis system (Spark Holland, Netherlands) and the Hysphere C18 HD cartridge (2x10 mm) (Spark Holland, Aj Emmen, Netherlands), and analytes were eluted during the mobile phase. Separation was performed with Synergy fusion-RP analytical column (250 mm x 2.0 mm, 4 µm particle diameter) (Phenomenex, Torrance, CA, USA) preceded by a guard column (4 mm x 2 mm) of the same packing material from Phenomenex. The mobile phase was a gradient of a mixture of 5 mM ammonium formate/formic acid 0.01% and acetonitrile/formic acid 0.01%. The flow rate was 0.2 mL/min, and the temperature 35°C. LC/MSMS analyses were performed with a system composed of a Waters alliance 2690 LC pump equipped with an autosampler and connected in series with a Quattro Ultima triple-quadrupole mass spectrometer (Waters-Micromass, Manchester, UK), equipped for

electrospray ionization. Acquisition took place in multiple reaction monitoring mode, with two transitions per compound (one for quantification and one for confirmation) in positive ionization mode. All validation procedures were performed with fresh samples of herbicide-free human urine. The limit of detection (LOD) was defined as a signal three times the background noise at the lowest analyte concentration assayed, and the limit of quantification (LOQ) as ten times the background noise. LOQs ranged from 0.001 to 1.7 µg/L. Concentration ranges were linear from 0.010 to 10 µg/L for herbicides with the lowest LODs (metolachlor, desethylatrazine, alachlor, 2,6-diethylaniline) and were linear from the LOD for the others. Average recoveries were 100% +/- 20%, with coefficients of variation ranging from 0.1 to 13.9%. Stability tests showed, among doped human urinary samples, a slight decrease in levels after 32 hours at ambient temperatures (respectively -7% and -9% for atrazine and atrazine mercapturate) and no changes according to the presence of nitric acid. This method is reported to provide minimal sample handling, good extraction recovery, with satisfactory LODs and LOQs (Panuwet et al. 2008, 2010).

Definition of exposure profiles from urinary concentrations

Most human biomonitoring studies assess atrazine exposure by measuring atrazine mercapturate in urine samples. This metabolite is formed directly from atrazine and cannot result from other atrazine degradation processes in the environment or organisms (See Figure 1 in Barr et al. 2007). However, measurement of this metabolite alone misrepresents total atrazine exposure (Barr et al. 2007). In our study, mothers were considered exposed to atrazine when atrazine or at least one of its specific metabolites (i.e., formed only from atrazine degradation processes: atrazine mercapturate, desethylatrazine, hydroxyatrazine, or hydroxydesethylatrazine) was quantified in their urine. Among them, those with quantifiable levels of atrazine or atrazine mercapturate were classified as “directly exposed” to atrazine.

We classified mothers as unexposed if there was no trace (i.e., <LOQ) of atrazine or any of its four specific metabolites in their urine sample. We used a similar strategy to classify exposure to simazine and its specific metabolites. We also classified women as exposed to dealkylated or hydroxylated triazine metabolites if at least one metabolite in the respective group was quantified in the urine sample, and classified women as exposed to alachlor if alachlor or its metabolite 2,6-diethylaniline was quantified. Molar concentrations (nmol/L) were summed to compute combined biomarker levels for groups of related compounds. For this computation, non-quantified values were replaced by the LOD divided by the square root of two, and molar concentrations were computed by dividing the chemical determination value, expressed in $\mu\text{g/L}$, by the molar mass.

Statistical analyses

Logistic models were used to estimate associations between biomarkers and adverse birth outcomes. We did not incorporate case-cohort sampling probabilities because the case-cohort odds ratio approximates the case-control odds ratio when the outcome is rare (Ahrens and Pigeot 2004). Covariates considered in multivariate analyses were known or suspected risk factors and factors that predicted outcomes or urinary concentrations in this cohort. An initial list of covariates, used for the FGR models, included: year of enrollment (2002-2003, 2004, 2005-2006), residence district (Ille-et-Vilaine, Côtes d'Armor, Finistère or other), maternal age (<25, 25-30, 30-35, >35 years old), education level (primary/secondary school, baccalaureate, technical school/post-graduate), smoking status at enrollment (nonsmoker, nonsmoker who smoked at conception, smoker), alcohol consumption at enrollment (abstinent, occasional, ≥ 1 drink per day), high blood pressure before/during pregnancy, season at conception, thawing and re-freezing process (that concerned a subset of 52 samples), and urine creatinine level (in g/L). For SHC models, we also considered cesarean delivery

(since passage through the birth canal may influence head circumference at birth), parity (1 if the current pregnancy is the first, 2, ≥ 3), pre-pregnancy body mass index (BMI) (<18.5 , 18.5-25, 25-30, ≥ 30 kg/m²). For congenital anomalies models, we considered additionally to the initial list, folic acid intake during the three months before enrollment (through vitamin supplementation), maternal fever in early pregnancy (prior to enrollment), child's sex, parity, pre-pregnancy BMI, mother's occupational exposure to solvents (no; yes; no job) (in accordance with previous associations observed from this cohort, Garlantézec et al. 2009), and gestational age at birth (in weeks). Linear models were used to estimate associations between biomarkers and birth weight, birth length and head circumference as continuous outcomes, using the same initial list of covariates plus child's sex, parity, pre-pregnancy BMI, gestational age at birth and squared gestational age, as well as fish intake (3 categories) and shellfish intake (3 categories) according to previous results from this cohort (Guldner et al. 2007), and, for head circumference, cesarean delivery. For all outcomes, the list of covariates also included exposures to herbicides, as categorical variable as defined above.

Final models were obtained using a backward selection process of the covariates ($p < 0.20$) and forcing in models the exposure of interest. Additional analyses considered biomarker levels as continuous in models and were thus restricted to samples with quantifiable levels of the compound of interest. SAS software was used for all calculations (SAS/STAT version 9.1, SAS Institute, Inc., Cary, NC).

Results

Population

Table 1 describes the case-cohort population. Most women in the random subcohort were recruited in Ille-et-Vilaine during 2003 and 2004. Brittany's population density is low, and

18% of the women lived in cities (>20,000 inhabitants). Average maternal age was 30 years, and most were postgraduate. Few (n=8) were farmers. At enrollment, mean gestational age was 11.7 weeks, 16% of mothers smoked, and 15% reported drinking alcohol occasionally or regularly. Pre-pregnancy BMI was 18.5-25 kg/m² (normal range) for 75%. Five percent had high blood pressure during pregnancy, and 17% of deliveries were by cesarean.

Major congenital anomalies were less frequent for women who conceived during the summer (compared to spring, OR=0.5; 95% CI: 0.2-1.0), and for those with periconceptual folic acid supplementation (OR=0.4; 95% CI: 0.2-0.9). Risk factors for FGR were low maternal education level (OR primary/secondary school versus baccalaureate=1.7; 95%CI: 1.1-2.9), smoking (OR=1.7; 95% CI: 1.1-2.5) or alcohol consumption (OR >1drink/day versus abstinent=2.5; 95% CI: 0.9-6.9) in early pregnancy, and high blood pressure during pregnancy (OR=2.9; 95%CI: 1.7-5.1). SHC was less common in the babies of women living in cities (OR cities versus rural or semi/rural areas=0.5; 95%CI: 0.3-1.0), who were multiparous (OR_{≥3} versus 1=0.5; 95%CI: 0.2-0.9), or who had cesarean versus vaginal deliveries (OR=0.5; 95%CI: 0.3-1.0), and SHC was more common in boys than girls (OR=3.4; 95%CI: 2.1-5.7).

Urinary measurements of atrazine and other herbicides

The 579 subcohort urine samples included only 10 (2%) with quantifiable atrazine and 24 (4%) with quantifiable atrazine mercapturate (Table 2); the prevalence of direct exposure to atrazine (quantification of at least one of these two compounds) was 6% (32 samples). Quantifiable levels of simazine or simazine mercapturate were observed in 8% of the subcohort. Dealkylated triazine metabolites were quantified in 20% of subcohort urine samples and hydroxylated triazine metabolites in 40%. The highest urinary concentrations

were observed among the dealkylated triazine metabolites. Median and maximum values were lowest for the chloroacetanilide compounds and for atrazine mercapturate.

Factors potentially associated with quantification of urinary atrazine or other herbicides

Evidence of direct exposure to atrazine (i.e., quantification of atrazine or atrazine mercapturate) was observed in samples collected during all years of the study (Figure 1). The proportion of samples with evidence of direct exposure did not differ substantially between samples from women living in cities and rural areas, or according to categories of estimated atrazine-contaminated tap water consumption. It was, however, higher in samples from women with the highest fish intake. We observed similar conclusions for alachlor exposure.

Simazine or simazine mercapturate, and dealkylated and hydroxylated triazine metabolites were quantified more frequently in urine samples of women living in the Côtes d'Armor district than in other municipalities, in rural areas than in cities, and in municipalities with the highest level of atrazine-contaminated tap water. The frequency of the dealkylated metabolites in urine decreased with the proportion of land devoted to corn in the mothers' home municipality.

Chloroacetanilide compounds were quantified more frequently in samples from women residing in Côtes d'Armor than women in other districts. Living in rural areas or near corn crops did not appear to explain the presence of metolachlor in urine samples, but the proportion of samples with evidence of metolachlor exposure was higher in samples from women with the highest intakes of fish and seafood.

In addition, our data showed no correlation between concentrations of triazines and gestational age at urine collection (4-19 weeks) and no relation between the biomarker levels and the time (in days) that samples spent at ambient temperatures (results not shown),

suggesting that it is unlikely that these sources of between-subject variation affected our results.

Association between exposure to atrazine and to other herbicides and adverse pregnancy outcomes

Table 3 describes the herbicide compounds that co-occurred with atrazine exposure in this study. In the 32 urine samples from women in the subcohort with evidence of direct exposure to atrazine, we also quantified alachlor, metolachlor, hydroxyatrazine, acetochlor, simazine, simazine mercapturate, desisopropyl desethylatrazine and its hydroxylated form, more frequently than expected: alachlor was found in 14 samples (44% compared with 10% in the subcohort), metolachlor in 16 samples (50% compared with 25%), and hydroxyatrazine in 10 samples (31% compared with 10%).

Major congenital anomalies were not associated with biomarkers of triazine exposure, but were associated with exposure to simazine (OR 1.8; 95% CI: 1.0-3.5, $p=0.07$) (Table 4). Atrazine exposure (i.e., quantification of atrazine or one of its specific metabolites) was associated with FGR (OR=1.5; 95% CI: 1.0-2.2) and SHC (OR=1.7; 95% CI: 1.0-2.7). Associations with direct exposure to atrazine were also positive but were not statistically significant (Table 4). There was no evidence of an association between exposure to other triazines and FGR or SHC, or between chloroacetanilide herbicides and any of the dichotomous pregnancy outcomes.

Birth weight, birth length, and head circumference were all lower in babies of women with atrazine or atrazine mercapturate quantified in their urine than in babies of women not exposed to atrazine, the decrease being statistically significant for birth weight (-151 g, 95% CI -293 to -9) (Table 5). Growth indicators were positively associated with direct exposure to simazine (birth weight +130 g, 95% CI: 5 to 255; length +0.61 cm, 95% CI: 0.1 to 1.2; head

circumference +0.38 cm, 95% CI: -0.03 to 0.8) and with alachlor or its metabolite in urine (for birth weight +91 g, 95% CI: 3 to 179).

Analyses of the babies of the 144 women with quantifiable metolachlor indicated a significant decrease in head circumference (beta=-0.53 cm, 95% CI -0.96 to -0.10 for a 1-nmol/L increase in metolachlor) and non-significant decreases in birth weight (-21 g, 95% CI -151 to 109) and birth length (-0.24 cm, 95% CI -0.78 to 0.31).

Additional analyses showed a weak positive association between quantification of atrazine or one of its specific metabolites in urine and male genital anomalies (OR 1.4, 95% CI: 0.6-3.2 based on 5 cases classified as exposed and 18 as unexposed) and a stronger but non-significant association with evidence of direct atrazine exposure (OR 2.3, 95% CI: 0.6-8.4 based on 3 cases classified as exposed and 18 as unexposed) compared to no exposure. Among the 35 mothers with quantified atrazine or atrazine mercapturate in their urine, the OR for male genital anomalies associated with a 1-nmol/L increase in these compounds was 2.8 (95% CI: 0.9-8.6).

Discussion

This study showed the presence of atrazine residues in urine samples of pregnant women up to three years after atrazine was banned in Europe. Only a few studies have measured urinary levels of triazine compounds in general and nonfarm populations (Curwin et al. 2007, Bakke et al. 2009), mainly in the United States; none focused on pregnant women. Curwin et al. (2007) studied 51 nonfarm children in Iowa in 2001 and observed higher levels of atrazine mercapturate than in our study: 12% had detectable levels (and their LOD at 1.16 µg/L was higher) and the maximum level was 2.2 µg/L. Another Iowa study of 10 nonfarm men in 2002-2003 observed urinary traces of atrazine mercapturate similar to our study (6% of detectable levels, with an LOD below 0.2 µg/L) (Bakke et al. 2009). As Barr et al. (2007)

suggested and our results confirm, measurement of atrazine mercapturate alone misrepresents and underestimates total atrazine exposure. We measured urinary concentrations of other triazine metabolites that may also be environmental degradates and found higher detection frequencies and levels for most of them than for atrazine or atrazine mercapturate. That metabolite traces are more frequent than the parent substance has also been generally observed in environmental media. The detection of atrazine and simazine residues in urine samples of pregnant women up to three years after they were banned in Europe in 2003 suggests that these pesticides persist several years in the environment. We however cannot rule out the possibility of uses on crops of forbidden products that represented 6% of the used products in 2006 in Brittany (AGRESTE Bretagne 2009). In the present study, there was no evidence of a link between herbicide exposure biomarkers and area devoted to corn production in the mother's municipality of residence. Although our assessment of atrazine contamination in tap water was crude, limited to the Ille-et-Vilaine district and to a period earlier than the study period, our descriptive results suggest that water contamination was a likely determinant of urinary levels of triazine compounds in our Britton population. Curwin et al. (2007) found high correlations between the atrazine mercapturate urinary levels of members of the same family in nonfarm families and suggested possible common sources of exposure for them, such as diet or water following environmental atrazine contamination.

Our study suggests that environmental atrazine exposure may impair fetal growth, as shown by outcomes such as FGR and SHC that may be considered as pathological entities. This result was reinforced by inverse associations between atrazine exposure and the crude fetal growth indicators, especially birth weight. To our knowledge, until now only limited animal data have suggested that gestational atrazine exposure may affect fetal growth. Although Rayner et al. (2004) showed a decrease in rat pup weight after *in utero* exposure to atrazine

alone, the same team observed the opposite effect when dams were exposed to a low-level mixture of atrazine and four degradates (Enoch et al. 2007). Little is known about the potential developmental toxicity of either the dealkylated or hydroxylated forms of triazine compounds. Other studies have suggested that dealkylated atrazine metabolites that retain the chlorine molecule may be as potent as the parent substance, whereas hydroxylated metabolites have lower potency (Stoker et al. 2002, Laws et al. 2003). Our study suggests that any adverse effect of these dealkylated and hydroxylated compounds on fetal growth outcomes would likely be driven mainly by metabolites specific to atrazine (desethylatrazine, hydroxyatrazine, or hydroxydesethylatrazine) and not by other dealkylated or hydroxylated triazine metabolites. Finally, our results are consistent with those of Munger et al. (1997), Villanueva et al. (2005), and Ochoa-Acuña et al. (2009), all of whom examined exposure during pregnancy to atrazine and its degradates by consumption of tap water. We did not, however, confirm associations between major congenital anomalies and atrazine exposure suggested in several recent ecological studies (Davis 2005, Mattix et al. 2007, Winchester et al. 2009, Waller et al. 2010). A meta-analysis has already linked hypospadias to parental pesticide exposure, with possible endocrine-mediated effects (Rocheleau et al. 2009). We found only one mammal experiment that observed hypospadias in rats treated with atrazine (Wu et al. 2007). Our finding of an association between male genital anomalies and atrazine or atrazine mercapturate levels in early-pregnancy urine samples was based on small numbers, and requires replication. It would reinforce the hypothesis of endocrine-mediated effects of low levels of atrazine exposure, formulated from animal observations in laboratories and in the wild (Hayes et al. 2006).

To our knowledge, our study suggests, for the first time, that simazine exposure and alachlor exposure may be positively associated with fetal growth indicators. These results need however to be replicated. Finally, our study suggests that metolachlor exposure levels may be

inversely correlated with head circumference, consistent with the recent report by Barr et al. (2009) of an inverse correlation with birth weight. In the present study, co-exposures to herbicides were taken into account as potential confounding factors in all models.

Because cohort members were asked to provide only one urine sample for this study, possible intraindividual variability cannot be considered. We assumed that one sample was enough to reflect chronic exposure resulting from environmental atrazine contamination. Chemical stability analyses suggest that our study may have non-differentially underestimated atrazine exposure levels, due to routine post shipment of urine samples.

Conclusion

This is the first study to assess the association between urinary biomarkers of exposure to triazine compounds during pregnancy and adverse birth outcomes and to take into account exposure to mixtures of atrazine, its environmental degradates, and other co-occurring herbicides. Because atrazine is banned in Europe, this study deals with environmental contamination by atrazine and its persistence in environment, which leads to low levels of atrazine exposure. Our results thus raise particular concerns for countries where atrazine is still in use.

References

AGRESTE Bretagne. Ministère de l'alimentation, de l'agriculture et de la pêche. 2001.

Enquêtes sur les pratiques culturelles en 2001 [in French]. Available:

http://draaf.bretagne.agriculture.gouv.fr/corpep/IMG/pdf/Synthese_Mol_internet.pdf

[accessed 10 December 2009]

AGRESTE Bretagne. Ministère de l'alimentation, de l'agriculture et de la pêche. 2009.

Enquêtes sur les pratiques culturelles en 2006, les pratiques phytosanitaires progressent avec la réglementation [in French]. Available:

http://www.agreste.agriculture.gouv.fr/IMG/pdf_R5309A06.pdf [accessed 6 January 2010]

Ahrens W, Pigeot I eds. Handbook of Epidemiology. Chapter I.7 Modern Epidemiologic Study Designs. Springer, 2004.

Bakke B, De Roos AJ, Barr DB, Stewart PA, Blair A, Freeman LB et al. 2009. Exposure to atrazine and selected non-persistent pesticides among corn farmers during a growing season. *J Expo Sci Environ Epidemiol* 19:544-554.

Barr DB, Ananth CV, Yan X, Lashley S, Smulian JC, Ledoux TA et al. 2010. Pesticide concentrations in maternal and umbilical cord sera and their relation to birth outcomes in a population of pregnant women and newborns in New Jersey. *Sci Total Environ* 408:790-795.

Barr DB, Panuwet P, Nguyen JV, Udunka S, Needham LL. 2007. Assessing exposure to atrazine and its metabolites using biomonitoring. *Environ Health Perspect* 115:1474-1478.

Barr DB, Wang RY, Needham LL. 2005. Biologic monitoring of exposure to environmental chemicals throughout the life stages: requirements and issues for consideration for the National Children's Study. *Environ Health Perspect* 113(8):1083-1091

Benoît M, Bonicelli B, Guichard L, Delorme R, Faloya V, Ruelle B. 2005. Chapitre 2: Connaissance de l'utilisation des pesticides. In: Expertise scientifique collective Inra et

Cemagref "Pesticides, agriculture et environnement": Réduire l'utilisation des pesticides et en limiter les impacts environnementaux [in French].

Curwin BD, Hein MJ, Sanderson WT, Striley C, Heederik D, Kromhout H et al. 2007.

Urinary pesticide concentrations among children, mothers and fathers living in farm and non-farm households in Iowa. *Ann Occup Hyg* 51:53-65.

Davis JA. 2005. Environmental Atrazine Exposure and Congenital Malformations in New York State: An Ecologic Study. Abstract in: Teratology Society 45th Annual Meeting St. Pete Beach, Florida.

Enoch RR, Stanko JP, Greiner SN, Youngblood GL, Rayner JL, Fenton SE. 2007. Mammary gland development as a sensitive end point after acute prenatal exposure to an atrazine metabolite mixture in female Long-Evans rats. *Environ Health Perspect* 115:541-547.

Garlantézec R, Monfort C, Rouget F, Cordier S. 2009. Maternal occupational exposure to solvents and congenital malformations: a prospective study in the general population. *Occup Environ Med* 66:456-463.

Guldner L, Monfort C, Rouget F, Garlantézec R, Cordier S. 2007. Maternal fish and shellfish intake and pregnancy outcomes: a prospective cohort study in Brittany, France. *Environ Health* 24:6:33.

Hayes TB, Stuart AA, Mendoza M, Collins A, Noriega N, Vonk A et al. 2006.

Characterization of atrazine-induced gonadal malformations in African clawed frogs (*Xenopus laevis*) and comparisons with effects of an androgen antagonist (cyproterone acetate) and exogenous estrogen (17beta-estradiol): Support for the demasculinization/feminization hypothesis. *Environ Health Perspect* 114 Suppl 1:134-141.

Laws SC, Ferrell JM, Stoker TE, Cooper RL. 2003. Pubertal development in female Wistar rats following exposure to propazine and atrazine biotransformation by-products, diamino-S-chlorotriazine and hydroxyatrazine. *Toxicol Sci* 76:190-200.

- LeBaron H, MacFarland J, Burnside O, eds. 2008. The Triazine Herbicides 50 Years Revolutionizing Agriculture. p36. Available:
<http://www.elsevierdirect.com/ISBN/9780444511676/The-Triazine-Herbicides> [Accessed 10 December 2009]
- Mamelle N, Cochet V, Claris O. 2001. Definition of fetal growth restriction according to constitutional growth potential. *Biol Neonate* 80:277-285.
- Mamelle N, Munoz F, Grandjean H. 1996. Fetal growth from the AUDIPOG study. I. Establishment of reference curves [in French]. *J Gynecol Obstet Biol Reprod (Paris)* 25:61-70.
- Marchand D. Service Santé-Environnement. DRASS de Bretagne. 2006. Le contrôle sanitaire des pesticides dans les eaux destinées à l'alimentation humaine en 2005 [in French]. Available:
http://draaf.bretagne.agriculture.gouv.fr/corpep/IMG/pdf/DRASS_pesticides_2005_CORP_EP.pdf [Accessed 6 January 2010]
- Mattix KD, Winchester PD, Scherer LR. 2007. Incidence of abdominal wall defects is related to surface water atrazine and nitrate levels. *J Pediatr Surg* 42:947-949.
- Munger R, Isacson P, Hu S, Burns T, Hanson J, Lynch CF et al. 1997. Intrauterine growth retardation in Iowa communities with herbicide-contaminated drinking water supplies. *Environ Health Perspect* 105:308-314.
- Ochoa-Acuña H, Frankenberger J, Hahn L, Carbajo C. 2009. Drinking-water herbicide exposure in Indiana and prevalence of small-for-gestational-age and preterm delivery. *Environ Health Perspect* 117:1619-1624.
- Panuwet P, Nguyen JV, Kuklennyik P, Udunka SO, Needham LL, Barr DB. 2008. Quantification of atrazine and its metabolites in urine by on-line solid-phase extraction-

high-performance liquid chromatography-tandem mass spectrometry. *Anal Bioanal Chem* 391:1931-1939.

Panuwet P, Restrepo PA, Magsumbol M, Jung KY, Montesano MA, Needham LL, Barr DB. 2010. An improved high-performance liquid chromatography-tandem mass spectrometric method to measure atrazine and its metabolites in human urine. *J Chromatogr B Analyt Technol Biomed Life Sci* 878:957-962.

Rayner JL, Wood C, Fenton SE. 2004. Exposure parameters necessary for delayed puberty and mammary gland development in Long-Evans rats exposed in utero to atrazine. *Toxicol Appl Pharmacol* 195:23-34.

Réseau CORPEP. Direction régionale de l'environnement de Bretagne. 2008. Les pesticides dans les eaux superficielles bretonnes [in French]. Available: http://draf.bretagne.agriculture.gouv.fr/corpep/IMG/pdf/bilan_reseau_corpep_2007.pdf [accessed 6 January 2010]

Rocheleau CM, Romitti PA, Dennis LK. 2009. Pesticides and hypospadias: a meta-analysis. *J Pediatr Urol.* 5:17-24.

Stoker TE, Guidici DL, Laws SC, Cooper RL. 2002. The effects of atrazine metabolites on puberty and thyroid function in the male Wistar rat. *Toxicol Sci* 67:198-206.

Villanueva CM, Durand G, Coutté MB, Chevrier C, Cordier S. 2005. Atrazine in municipal drinking water and risk of low birth weight, preterm delivery, and small-for-gestational-age status. *Occup Environ Med* 62:400-405.

Waller SA, Paul K, Peterson SE, Hitti JE. 2010. Agricultural-related chemical exposures, season of conception, and risk of gastroschisis in Washington State. *Am J Obstet Gynecol* 202:241.

Winchester PD, Huskins J, Ying J. 2009. Agrichemicals in surface water and birth defects in the United States. *Acta Paediatr* 98:664-669.

Wu YG, Li SK, Xin ZC, Wang YS, Shou KR, Gao H et al. 2007. The establishment of hypospadias rat model and embryoteratogenic test of Atrazine [in Chinese, abstract only]. Zhonghua Zheng Xing Wai Ke Za Zhi 23:340-343.

Table 1: Characteristics of the case-cohort population of pregnant women assessed for urinary biomarkers of pesticide exposure (France, 2002-2006)

	Subcohort		Major congenital anomalies		FGR		SHC		
	n	%	n	%	n	%	n	%	
All	579		88		178		103		
Year of inclusion	2002-2003	292	50%	51	58%	89	50%	43	42%
	2004	206	36%	23	26%	60	34%	48	47%
	2005-2006	81	14%	14	16%	29	16%	12	12%
District	Ille-et-Vilaine	361	62%	63	72%	112	63%	58	56%
	Côtes d'Armor	158	27%	21	24%	53	30%	42	41%
	Finistère	35	6%	3	3%	8	4%	1	1%
	Other	25	4%	1	1%	5	3%	2	2%
Habitat	Rural/Semi-rural area (<20,000 inhabitants)	473	82%	72	82%	151	85%	92	89%
	Urban (≥20,000 inhabitants)	106	18%	16	18%	27	15%	11	11%
Maternal age (years)	<25	73	13%	9	10%	25	14%	16	15%
	25-30	224	39%	30	34%	60	34%	38	37%
	30-35	202	35%	34	39%	68	38%	39	38%
	≥35	80	14%	15	17%	25	14%	10	10%
	<i>Mean (SE) ; Min-max</i>	<i>30 (0.2); 19-44</i>							
Maternal education	Primary/secondary school	109	19%	17	19%	47	26%	15	15%
	Baccalaureate	109	19%	16	18%	28	16%	24	23%
	Technical school/Post-secondary	360	62%	55	63%	103	58%	64	62%
	Missing	1	-						
Maternal farm work	Yes	8	1.7%	1	1.1%	3	2%	2	2%
Maternal smoking status at enrollment	Nonsmoker	411	72%	70	80%	118	66%	66	65%
	Nonsmoker but smoked at conception	70	12%	9	10%	17	10%	13	13%
	Smoker	91	16%	9	10%	43	24%	23	22%
	Missing	7	-					1	-
Maternal drinking status at enrollment	Abstinent	486	85%	77	89%	149	85%	89	89%
	Occasional (<one drink per day)	75	13%	9	10%	19	11%	8	8%
	≥1 drink per day	9	2%	1	1%	7	4%	3	3%
	Missing	9	-	1	-	3	-	3	-
Season of conception	Spring	121	21%	23	26%	35	20%	23	22%
	Summer	162	28%	15	17%	48	27%	34	33%
	Autumn	142	24%	29	33%	45	25%	22	21%
	Winter	154	27%	21	24%	50	28%	24	23%
Pre-pregnancy body mass index (kg/m ²)	<18.5	51	9%	10	11%	12	7%	11	11%
	18.5-25	426	74%	68	77%	133	75%	77	75%
	25-30	68	12%	6	7%	20	11%	8	8%
	≥30	27	5%	4	5%	13	7%	6	6%
	Missing	7	-					1	-
	<i>Mean (SE) ; Min-max</i>	<i>22.2 (0.2); 15-44</i>							
Parity	1	264	46%	43	49%	83	47%	57	55%
	2	207	36%	25	28%	70	39%	35	34%
	≥3	107	18%	20	23%	25	14%	11	11%
	Missing	1							
Folic acid intake (3 months before inclusion)	Yes	100	17%	7	8%	22	12%	16	16%
High blood pressure before/during pregnancy	Yes	32	6%	2	2%	26	15%	4	4%
Cesarean delivery	Yes	100	17%	14	16%	42	24%	10	10%
Gestational age at inclusion ^a	<i>Mean (SE); Min-max</i>	<i>11.7 (0.1); 4-19</i>							
Gestational duration ^a	<i>Mean (SE); Min-max</i>	<i>39.3 (0.07); 26-42</i>							
Infant characteristics:									
Sex	Girls	280	48%	35	42%	91	51%	22	21%
	Boys	299	52%	48	58%	87	49%	81	79%
	Missing			5 ^b					
Birth weight (g)	<i>Mean (SE); Min-max</i>	<i>3376 (20); 1050-4760</i>							
Birth length (cm)	<i>Mean (SE); Min-max</i>	<i>49.7 (0.1); 35-56</i>							
Head circumference at birth (cm)	<i>Mean (SE); Min-max</i>	<i>34.6 (0.07); 25-39</i>							

Abbreviations: FGR: fetal growth restriction; SHC: small head circumference.

^a In weeks of amenorrhea ^b Five medical pregnancy terminations without available data on fetus gender

Table 2: Levels of triazine and chloroacetanilide herbicides quantified in urine from 579 randomly selected women from the PELAGIE cohort in early pregnancy (France, 2002-2006)

	LOQ ($\mu\text{g/L}$)	n (%) >LOQ	Median (p95, Max) among quantified values
Triazine compounds (in $\mu\text{g/L}$):			
Atrazine	0.05	10 (2%)	0.12 (0.52 , 0.52)
Atrazine mercapturate	0.02	24 (4%)	0.05 (0.49 , 0.68)
Simazine	0.2	6 (1%)	1.0 (4.4 , 4.4)
Simazine mercapturate	0.06	44 (8%)	0.5 (3.4 , 4.6)
Desethylatrazine ^a	0.003	60 (10%)	0.1 (7.8 , 14.0)
Desisopropyl atrazine ^a	0.9	25 (4%)	1.4 (7.1 , 8.2)
2-chlorodiaminoatrazine ^a	0.5	42 (7%)	1.0 (2.9 , 5.0)
Hydroxyatrazine ^b	0.02	57 (10%)	0.1 (1.1 , 4.0)
Hydroxydesethylatrazine ^b	0.315	31 (5%)	0.6 (2.5 , 2.5)
Hydroxysimazine ^b	0.02	50 (9%)	0.08 (1.1 , 1.6)
Hydroxy desisopropyl atrazine ^b	0.15	90 (16%)	0.6 (3.5 , 4.5)
Ammeline ^b	0.25	79 (14%)	0.9 (5.9 , 7.1)
Chloroacetanilide compounds (in $\mu\text{g/L}$):			
Alachlor	0.006	64 (11%)	0.03 (0.21 , 0.65)
2,6-diethylaniline	0.01	54 (9%)	0.03 (0.13 , 0.82)
Metolachlor	0.002	144 (25%)	0.01 (0.18 , 1.14)
Acetochlor	0.02	51 (9%)	0.05 (0.17 , 1.04)
Creatinine (in $\mu\text{g/L}$):		579 (100%)	1009 (2014 , 3511)
Sum of compounds (in nmol/L)^c:			
Σ (Atrazine, Atrazine mercapturate)		32 (6%)	0.2 (1.4 , 4.4)
Σ (Simazine, Simazine mercapturate)		47 (8%)	2.0 (12.6 , 21.8)
Σ Dealkylated triazine metabolites		117 (20%)	5.3 (34.6 , 74.8)
Σ Hydroxylated triazine metabolites		229 (40%)	4.4 (30.9 , 66.5)
Σ (Alachlor, 2,6-diethylaniline)		103 (18%)	0.2 (1.0 , 7.2)

Abbreviations : LOQ: quantification limit; p95: 95th percentile

^a Dealkylated triazine metabolites (desethylatrazine, desisopropyl atrazine, 2-chlorodiaminoatrazine)

^b Hydroxylated triazine metabolites (hydroxyatrazine, hydroxysimazine, hydroxydesethylatrazine, hydroxy desisopropyl atrazine, ammeline)

^c Compounds were summed according to the definition of exposure profiles of interest, as stated in the Materials and Methods section. Example: Σ (Atrazine, Atrazine mercapturate) means that atrazine or atrazine mercapturate were found in 32 urine samples (sample may contain both compounds) with a median concentration of 0.2 nmol/L.

Table 3: Herbicide compounds co-occurring with atrazine exposure in urine samples of 579 pregnant women (France, 2002-2006)

Co-occurring compounds:	Numbers of urine samples with the co-occurring compound		
	Among samples with no exposure to atrazine ^a	Among samples with atrazine exposure ^b	Among samples with direct exposure to atrazine
(Total)	(425)	(154)	(32)
Atrazine	0	10	10
Atrazine mercapturate	0	24	24
Desethylatrazine	0	60	3
Hydroxyatrazine	0	57	10
Hydroxydesethylatrazine	0	31	3
Simazine	0	6	2
Simazine mercapturate	20	24	5
Hydroxysimazine	20	30	3
Desisopropyl atrazine	20	5	0
2-chlorodiaminoatrazine	27	15	5
Hydroxy desisopropyl atrazine	68	22	3
Ammeline	46	33	7
Alachlor	35	29	14
2,6-diethylaniline	39	15	6
Metolachlor	81	63	16
Acetolachlor	34	17	9

^a No quantified level of atrazine and its four specific metabolites (Desethylatrazine, hydroxyatrazine, hydroxydesethylatrazine and atrazine mercapturate).

^b At least one quantified compound among atrazine or its four specific metabolites

^c Quantified level of atrazine or atrazine mercapturate

Table 4: Association between pregnancy outcomes and exposure to herbicides assessed from urinary metabolites quantified in maternal samples collected in early pregnancy, adjusted for co-occurring exposures to other herbicides (France, 2002-2006)

	Groups of urinary metabolites quantified in urine samples	Major congenital anomalies			FGR			SHC		
		n	OR ^f	(95% CI)	n	OR ^g	(95% CI)	n	OR ^h	(95% CI)
Atrazine and its specific metabolites ^a	None	62	ref		117	ref		65	ref	
	At least one	26	1.2	(0.7, 2.1)	61	1.5	(1.0, 2.2) ^e	38	1.7	(1.0, 2.7) ^e
	At least atrazine or atrazine mercapturate (Direct exposure)	6	1.0	(0.3, 3.1)	14	1.6	(0.8, 3.1)	6	1.3	(0.5, 3.3)
Simazine and its specific metabolites ^b	None	71	ref		150	ref		90	ref	
	At least one	17	1.8	(1.0, 3.5)	28	1.1	(0.7, 1.8)	13	0.8	(0.4, 1.7)
	At least simazine or simazine mercapturate	9	1.6	(0.7, 3.8)	16	1.0	(0.5, 1.9)	7	0.6	(0.2, 1.6)
Dealkylated triazine metabolites ^c	No dealkylated metabolite	74	ref		140	ref		84	ref	
	At least one	14	0.9	(0.5, 1.7)	38	0.9	(0.6, 1.5)	19	0.8	(0.4, 1.4)
Hydroxylated triazine metabolites ^d	No hydroxylated metabolite	48	ref		107	ref		58	ref	
	At least one	40	1.3	(0.8, 2.2)	71	0.8	(0.6, 1.3)	45	1.1	(0.7, 1.7)
Alachlor	None	71	ref		144	ref		84	ref	
	At least alachlor or 2,6-diethylaniline	17	0.9	(0.4, 1.7)	34	1.3	(0.8, 2.0)	19	0.9	(0.5, 1.7)
Metolachlor	No	70	ref		133	ref		78	ref	
	Yes	18	0.8	(0.4, 1.4)	45	1.2	(0.8, 1.9)	25	1.0	(0.6, 1.7)
Acetochlor	No	82	ref		169	ref		94	ref	
	Yes	6	1.0	(0.4, 2.5)	9	0.6	(0.3, 1.2)	9	0.8	(0.4, 1.6)

Abbreviations: CI: confidence interval; FGR: fetal growth restriction; SHC: small head circumference for sex and gestational age
 Logistic models were adjusted according to backward selection (p<0.20). The exposure of interest was forced in the model (whatever its p-value). The retained covariates in final models are listed in ^f to ^h footnotes.

^a Desethylatrazine, hydroxyatrazine, hydroxydesethylatrazine and atrazine mercapturate. ^b Hydroxysimazine and simazine mercapturate.

^c Dealkylated metabolites: desethylatrazine, desisopropyl atrazine, 2-chlorodiaminoatrazine ^d Hydroxylated metabolites: hydroxyatrazine, hydroxydesethylatrazine, hydroxysimazine, hydroxy desisopropyl atrazine, ammeline ^e Statistical significance at 5% (the confidence interval does not include 1)

^f Retained covariates at p<0.20: year of enrollment, season at conception, mother's occupational exposure to solvents, gestational age at birth and simazine exposure (except for the models on dealkylated and hydroxylated metabolites)

^g Retained covariates at p<0.20: smoking status at enrollment, high blood pressure before/during pregnancy, thawing and re-freezing process, acetochlor exposure and atrazine exposure

^h Retained covariates at p<0.20: residence district, alcohol consumption at enrollment (except for the models on dealkylated and hydroxylated metabolites), thawing and re-freezing process, cesarean delivery, parity, atrazine exposure

Table 5: Association between intrauterine growth indicators and exposure to herbicides assessed from urinary metabolites quantified in maternal samples collected in early pregnancy, adjusted for co-occurring exposures to other herbicides (France, 2002-2006)

	Groups of urinary metabolites quantified in urine samples	Sub-cohort n	Birth weight (g)			Birth length (cm)			Head circumference (cm)		
			β^e	(95% CI)	<i>p</i>	β^f	(95% CI)	<i>p</i>	β^g	(95% CI)	<i>p</i>
Atrazine and its specific metabolites ^a	None	425	ref			ref			ref		
	At least one	154	-57	(-130, 17)	0.13	-0.22	(-0.5, 0.1)	0.20	-0.18	(-0.4, 0.1)	0.16
	At least atrazine or atrazine mercapturate (Direct exposure)	32	-151	(-293, -9)	0.04	-0.44	(-1.1, 0.2)	0.19	-0.39	(-0.9, 0.1)	0.10
Simazine and its specific metabolites ^b	None	506	ref			ref			ref		
	At least one	73	40	(-63, 142)	0.45	0.17	(-0.3, 0.6)	0.48	0.27	(-0.1, 0.6)	0.11
	At least simazine or simazine mercapturate	47	130	(5, 255)	0.04	0.61	(0.1, 1.2)	0.03	0.38	(-0.03, 0.8)	0.07
Dealkylated triazine metabolites ^c	No dealkylated metabolite	462	ref			ref			ref		
	At least one	117	-18	(-99, 62)	0.65	0.03	(-0.3, 0.4)	0.87	0.18	(-0.1, 0.4)	0.18
Hydroxylated triazine metabolites ^d	No hydroxylated metabolite	350	ref			ref			ref		
	At least one	229	-12	(-79, 55)	0.72	0.002	(-0.3, 0.3)	0.99	-0.09	(-0.3, 0.1)	0.41
Alachlor	None	476	ref			ref			ref		
	At least alachlor or 2,6-diethylaniline	103	91	(3, 179)	0.04	0.35	(-0.05, 0.7)	0.08	0.11	(-0.2, 0.4)	0.45
Metolachlor	No	435	ref			ref			ref		
	Yes	144	5	(-78, 87)	0.91	-0.20	(-0.6, 0.1)	0.25	0.04	(-0.2, 0.3)	0.73
Acetochlor	No	528	ref			ref			ref		
	Yes	51	65	(-53, 183)	0.28	0.13	(-0.4, 0.7)	0.63	0.09	(-0.3, 0.5)	0.65

Abbreviations: CI: confidence interval; FGR: fetal growth restriction; SHC: small head circumference for sex and gestational age

Linear models were adjusted according to backward selection ($p < 0.20$). The exposure of interest was forced in the model (whatever its *p*-value). The retained covariates in final models are listed in ^e to ^g footnotes.

^a Desethylatrazine, hydroxyatrazine, hydroxydesethylatrazine and atrazine mercapturate. ^b Hydroxysimazine and simazine mercapturate.

^c Dealkylated metabolites: desethylatrazine, desisopropyl atrazine, 2-chlorodiaminoatrazine ^d Hydroxylated metabolites: hydroxyatrazine, hydroxydesethylatrazine, hydroxysimazine, hydroxy desisopropyl atrazine, ammeline ^e Retained covariates: year of enrollment, education level, smoking status at enrollment, high blood pressure before/during pregnancy, thawing and re-freezing process, pre-pregnancy body mass index, child's sex, shellfish intake (except for the models on dealkylated and hydroxylated metabolites), gestational age at birth, squared gestational age, alachlor exposure and atrazine exposure

^f Retained covariates: year of enrollment, smoking status at enrollment, high blood pressure before/during pregnancy, season at conception, pre-pregnancy body mass index, child's sex, shellfish intake, gestational age at birth, squared gestational age, alachlor exposure and atrazine exposure (except for the models on dealkylated and hydroxylated metabolites and on metolachlor)

^g Retained covariates: residence district, smoking status at enrollment, pre-pregnancy body mass index, child's sex, cesarean delivery, parity, shellfish intake, gestational age at birth, squared gestational age, atrazine and simazine exposures (except for the models on dealkylated and hydroxylated metabolites)

Figure legend:

Figure 1

Dealkylated triazine metabolites: desethylatrazine, desisopropyl atrazine, 2-chlorodiaminoatrazine

Hydroxylated triazine metabolites: hydroxyatrazine, hydroxysimazine, hydroxydesethylatrazine, hydroxy desisopropyl atrazine, ammeline

